

Cluster Probability in Bootstrap Percolation

A. B. Harris and Andrea J. Liu

Department of Physics and Astronomy, University of Pennsylvania, Philadelphia, PA 19104

(Dated: September 14, 2007)

We develop a recursive formula for the probability of a k -cluster in bootstrap percolation.

PACS numbers: 64.60.Ak

I. INTRODUCTION

The model of bootstrap percolation¹ (or k -core percolation, as we will refer to it) has a connection to jamming transitions, including the glass transition.² Many features of dynamical arrest in glassforming liquids are captured by kinetically-constrained spin models,³ which map onto k -core percolation and its variants. In addition, Schwarz, et al.⁴ have argued that there is an analogy (but not a strict mapping) between k -core percolation and the jamming transition of frictionless granular packings^{5,6}, similar to the analogy between k -core percolation and rigidity percolation.⁷ In the mean-field, or infinite-dimensional limit, the k -core percolation transition was shown to be peculiar in Ref. 4 and subsequently in Ref. 8. The transition is discontinuous,^{1,9} but also exhibits power-law scaling^{1,9} and a diverging susceptibility and length scale⁴. This unusual type of transition has been called a random first-order phase transition.¹⁰ What makes the mean-field k -core percolation transition particularly interesting is that its exponents are identical to the mean-field exponents of several models that have been proposed for jamming transitions,^{10,11,12,13} raising the possibility that k -core percolation may be in the same universality class as these models.

For jamming transitions, the random first-order nature of the transition does not appear to be an artifact of the mean-field approximation. Numerical simulations^{5,6} and scaling arguments^{14,15} suggest that the jamming transition of frictionless granular packings (Point J) is discontinuous with a diverging length scale in two and three dimensions. It has also been argued that the finite-dimensional glass transition, if it exists as a true phase transition, has a mixed character, with a discontinuity in the infinite-time limit of the dynamical structure factor and a diverging dynamical length scale.^{16,17} This raises the question of the nature of the k -core percolation transition in finite dimensions d . Expansions in $1/d$ indicate¹⁸ that the mixed nature of the transition should persist for a range of large but finite d . In two dimensions, specific variants of k -core percolation have been identified that are rigorously known to exhibit discontinuous transitions with diverging length scales, both at $p = 1$ (full occupancy)^{13,19} and $p < 1$ (partial occupancy).¹⁷ Numerical studies of other variants⁴ also point to the possibility of mixed transitions in finite dimensions at $p < 1$. However, it is also known that some realizations of the model, such as $k = 3$ on the two-dimensional triangular lattice,

exhibit continuous transitions.²⁰ Thus, in finite dimensions, it appears that different versions of k -core models can exhibit at least three different types of transitions: continuous transitions, mixed transitions at $p = 1$ and mixed transitions at $p < 1$.

In order to sort through this confusing state of affairs, it would be useful to map k -core percolation onto a spin model, which could then be studied by a wide variety of methods. Such an approach proved extremely powerful for ordinary percolation. A necessary first step towards this goal is to be able to count k -core clusters with the correct probabilities. In this paper, we show how this can be done by deriving a relation for the probability distribution of clusters.

The k -core ensemble is defined as follows. We start with the ensemble of $2^{\mathcal{N}}$ percolation configurations on a *finite* lattice of \mathcal{N} bonds. Each percolation configuration \mathcal{C}_p has probability

$$P_p(\mathcal{C}_p) = p^{n_{\text{occ}}(\mathcal{C}_p)}(1-p)^{n_{\text{vac}}(\mathcal{C}_p)}, \quad (1)$$

where $n_{\text{occ}}(\mathcal{C}_p)$ and $n_{\text{vac}}(\mathcal{C}_p)$ are, respectively, the number of occupied and vacant bonds in the percolation configuration \mathcal{C}_p . A k -core configuration \mathcal{C}_k is obtained from \mathcal{C}_p by removing all bonds which intersect a site to which there are less than k remaining bonds. This process is recursively continued because as bonds are so removed, they may cause more sites to have less than k neighbors. After this culling, the remaining k -cluster is one in which all sites have at least k occupied bonds. The weight of the configuration \mathcal{C}_k is the sum of the percolation probabilities over all the percolation configurations which gave rise to \mathcal{C}_k .

In the percolation problem one can easily establish that the probability of finding an isolated cluster²¹ Γ_p is given by

$$P_p(\Gamma_p) = p^{n(\Gamma_p)}(1-p)^{t(\Gamma_p)}, \quad (2)$$

where $n(\Gamma_p)$ is the number of occupied bonds in the cluster Γ_p and $t(\Gamma_p)$ is the number of perimeter bonds which must be unoccupied in order to define the limit of the percolation cluster Γ_p . One would like to write a similar formula for $P_k(\Gamma)$, the probability of a k -cluster Γ_k , but it is obvious that this is not quite so trivial because of the complications of the culling process. It is the purpose of this paper to develop such a formula.

II. FORMULATION

To proceed we introduce the function $e_\Gamma(\mathcal{C}_p)$ which is defined as unity if the percolation configuration \mathcal{C}_p , when culled, gives rise to the rooted k -cluster Γ . A rooted cluster includes the site at the origin and two rooted clusters are distinct unless their list of occupied bonds coincides exactly. The final culled configuration must contain the rooted k -cluster Γ , but can also contain an arbitrary number of k -clusters disconnected from Γ . If culling of \mathcal{C}_p does not produce a configuration containing the rooted cluster Γ , then $e_\Gamma(\mathcal{C}_p)$ is zero. With this definition it seems obvious that

$$P_k(\Gamma) = \sum_{\mathcal{C}_p} e_\Gamma(\mathcal{C}_p) P_p(\mathcal{C}_p). \quad (3)$$

This relation expresses the fact that the cluster Γ inherits the total percolation probability of all configurations which contain the cluster Γ after culling. Note that a fixed Γ inherits probability from not only single clusters which, when culled, yield Γ , but also from the vast array of configurations which, when culled, yield Γ together with arbitrary disconnected k -clusters. Now it is clear that the sum over \mathcal{C}_p can be restricted to configurations which either are equal to Γ or include it as a proper subset. So we write

$$\begin{aligned} P_k(\Gamma) &= \sum_{\mathcal{C}_p: \mathcal{C}_p \geq \Gamma} e_\Gamma(\mathcal{C}_p) P_p(\mathcal{C}_p) \\ &= \sum_{\mathcal{C}_p: \mathcal{C}_p \geq \Gamma} \left(1 - [1 - e_\Gamma(\mathcal{C}_p)]\right) P_p(\mathcal{C}_p). \end{aligned} \quad (4)$$

But

$$\sum_{\mathcal{C}_p: \mathcal{C}_p \geq \Gamma} P_p(\mathcal{C}_p) = p^{n(\Gamma)} \quad (5)$$

because after we occupy the bonds of Γ (with probability $p^{n(\Gamma)}$) we sum over all states of all the bonds not in Γ (which are occupied with probability p and vacant with probability $1 - p$). Thus

$$P_k(\Gamma) = p^{n(\Gamma)} - \sum_{\mathcal{C}_p: \mathcal{C}_p \geq \Gamma} [1 - e_\Gamma(\mathcal{C}_p)] P_p(\mathcal{C}_p). \quad (6)$$

The result of Eq. (2) for the probability of a percolation cluster can be calculated in this same style. Note the meaning of the factor $[1 - e_\Gamma(\mathcal{C}_p)]$: it is an indicator function such that culling does *not* lead to the k -cluster Γ . But since \mathcal{C}_p contains Γ but does not cull to Γ , it must be that the configuration \mathcal{C}_p leads to a cluster $\Gamma_>$ which includes Γ as a proper subset. So we have

$$\begin{aligned} P_k(\Gamma) &= p^{n(\Gamma)} - \sum_{\Gamma': \Gamma' > \Gamma} \sum_{\mathcal{C}_p: \mathcal{C}_p \geq \Gamma'} e_{\Gamma'}(\mathcal{C}_p) P_p(\mathcal{C}_p) \\ &= p^{n(\Gamma)} - \sum_{\Gamma': \Gamma' > \Gamma} P_k(\Gamma'). \end{aligned} \quad (7)$$

Note that the sum over k -clusters Γ' does *not* include $\Gamma' = \Gamma$. Equation (7) is the principal result of this paper. On a qualitative level we can see the following: for percolation Eq. (2) gives what could be called a perimeter renormalization $(1 - p)^{n_i(C)}$, whereas the subtraction terms in Eq. (7) give a much weaker ‘‘culling renormalization.’’ In the extreme case of lattice animals, there is no perimeter renormalization at all. In that case $P(C) = p^{n(C)}$ (where p is interpreted to be the bond fugacity, usually denoted K). The continuous aspect of k -core percolation has⁴ the same anomalous correlation length exponent $\nu = 1/4$ as does lattice animals in infinite dimensions.²² A heuristic explanation of this is that the culling renormalization is probably closer to the absence of a perimeter renormalization in lattice animals than to the perimeter renormalization of percolation.

Now we briefly explore some consequences of the above result. We iterate Eq. (7) to get

$$\begin{aligned} P_k(\Gamma_k) &= p^{n(\Gamma_k)} - \sum_{\Gamma_k^{(1)} > \Gamma_k} p^{n(\Gamma_k^{(1)})} + \sum_{\Gamma_k^{(2)} > \Gamma_k^{(1)} > \Gamma_k} p^{n(\Gamma_k^{(2)})} \\ &\quad \dots + (-1)^m \sum_{\Gamma_k^{(m)} > \Gamma_k^{(m-1)} > \dots > \Gamma_k^{(1)} > \Gamma_k} p^{n(\Gamma_k^{(m)})} + \dots \\ &= p^{n(\Gamma_k)} + \sum_{m=1}^{m_{\max}} (-1)^m \sum_{\{\Gamma_k^{(m)}\}} p^{n(\Gamma_k^{(m)})}, \end{aligned} \quad (8)$$

where, for simplicity, we do not write out (in the last line) the inclusions. Because we are dealing with a finite lattice, the index m is bounded by a maximum value, denoted m_{\max} .

Next we consider the calculation of the connectedness susceptibility, χ , which can be written as

$$\chi = \sum_{\Gamma_k} P_k(\Gamma_k) n(\Gamma_k)^2. \quad (9)$$

Thus we write

$$\begin{aligned} \chi &= \sum_{\Gamma_k} n(\Gamma_k)^2 \left[p^{n(\Gamma_k)} \right. \\ &\quad \left. + \sum_{m=1}^{m_{\max}} (-1)^m \sum_{\{\Gamma_k^{(m)}\}} p^{n(\Gamma_k^{(m)})} \right]. \end{aligned} \quad (10)$$

By relabeling clusters, this can be rearranged to combine all terms with the same factor $p^{n(\Gamma)}$, in which case one has

$$\begin{aligned} \chi &= \sum_{\Gamma_k} p^{n(\Gamma_k)} \left[n(\Gamma)^2 - \sum_{\Gamma_k^{(1)} < \Gamma} [n(\Gamma_k^{(1)})]^2 \right. \\ &\quad \dots + (-1)^m \sum_{\Gamma_k^{(m)} < \Gamma_k^{(m-1)} < \dots < \Gamma_k^{(1)} < \Gamma} [n(\Gamma_k^{(m)})]^2 \\ &\quad \left. + \dots \right]. \end{aligned} \quad (11)$$

This result no longer requires that \mathcal{N} be finite because for any finite cluster Γ_k , the number of terms coming from included subclusters in the sum inside the square brackets is independent of system size. Thus this formula presumably can be used as long as $p < p_k$ and it may yield a possible route to an analytic approach to the calculation of χ for finite dimensional lattices.

The result Eq. (11) can be generalized to any observable $G(\Gamma_k)$ that can be defined for a k -cluster Γ_k . In that case, the ensemble-averaged value of G is

$$\langle G \rangle = \sum_{\Gamma_k} p^{n(\Gamma_k)} G_c(\Gamma_k), \quad (12)$$

where the cumulant function G_c is

$$G_c(\Gamma_k) = G(\Gamma_k) - \sum_{\Gamma_k^{(1)} < \Gamma_k} G(\Gamma_k^{(1)})$$

$$\dots + (-1)^m \sum_{\Gamma_k^{(m)} < \Gamma_k^{(m-1)} \dots < \Gamma_k^{(1)} < \Gamma_k} G(\Gamma_k^{(m)}) + \dots \quad (13)$$

or more simply,

$$G_c(\Gamma_k) = G(\Gamma_k) - \sum_{\Gamma_k^{(1)} < \Gamma_k} G_c(\Gamma_k^{(1)}). \quad (14)$$

This cumulant subtraction for the k -core percolation problem is much more difficult to handle than it is for ordinary percolation.

We thank Vincenzo Vitelli for stimulating discussions and a critical reading of the manuscript. This work was supported by NSF-DMR-0605044 (AJL).

-
- ¹ J. Chalupa, P. L. Leath, and G. R. Reich, *J. Phys. C* **12**, L31 (1979).
² A. J. Liu and S. R. Nagel, Eds. *Jamming and Rheology* (Taylor and Francis: New York, 2001).
³ F. Ritort and P. Sollich, *Advances in Physics*, **52**, 219 (2003).
⁴ J. M. Schwarz, A. J. Liu, and L. Chayes, *Europhys. Lett.* **73**, 560 (2006).
⁵ C. S. O'Hern, S. A. Langer, A. J. Liu and S. R. Nagel, *Phys. Rev. Lett.* **88**, 075507 (2002); C. S. O'Hern, L. E. Silbert, A. J. Liu, and S. R. Nagel, *Phys. Rev. E* **68**, 011306 (2003).
⁶ L. E. Silbert, A. J. Liu, and S. R. Nagel, *Phys. Rev. Lett.* **95**, 098301 (2005).
⁷ C. Moukarzel, P. M. Duxbury, and P. L. Leath, *Phys. Rev. E* **55**, 5800 (1997).
⁸ S. N. Dorogovtsev, A. V. Goltsev and J. F. F. Mendes, *Phys. Rev. Lett.* **96**, 040601 (2006).
⁹ B. Pittel, J. Spencer, and N. Wormald, *J. Comb. Th. Series B* **67**, 111 (1996).
¹⁰ T. R. Kirkpatrick and P. G. Wolynes, *Phys. Rev. B* **36**, 8552 (1986); T. R. Kirkpatrick and D. Thirumalai, *Phys. Rev. Lett.* **58**, 2091 (1987).
¹¹ D. Das, J. Kondev, B. Chakraborty, *Europhys. Lett.* **61**, 506 (2003).
¹² G. Biroli and J.-P. Bouchaud, *Europhys. Lett.* **67**, 21 (2004).
¹³ C. Toninelli, G. Biroli, D. S. Fisher, *Phys. Rev. Lett.* **92**, 185504 (2004).
¹⁴ M. Wyart, S. R. Nagel, T. A. Witten, *Europhys. Lett.* **72**, 486 (2005).
¹⁵ M. Wyart, L. E. Silbert, S. R. Nagel, T. A. Witten, *Phys. Rev. E* **72**, 051306 (2005).
¹⁶ T. R. Kirkpatrick, D. Thirumalai and P. G. Wolynes, *Phys. Rev. A*, **40**, 1045 (1989).
¹⁷ C. Toninelli, G. Biroli, D. S. Fisher, *Phys. Rev. Lett.* **96**, 035702 (2006).
¹⁸ A. B. Harris and J. Schwarz, *Phys. Rev. E* **72**, 046123 (2005).
¹⁹ P. De Gregorio, A. Lawlor, P. Bradley, K. A. Dawson, *Proc. Natl. Acad. Sci.* **102**, 5669 (2005).
²⁰ M. C. Medeiros and C. M. Chaves, *Physica A* **234**, 604 (1997); C. M. Chaves and B. Koiller, *Physica A* **218**, 271 (1995).
²¹ Single clusters are denoted by Γ with appropriate scripts. Configurations (which may consist of a union of clusters) are denoted \mathcal{C} with appropriate scripts.
²² T. C. Lubensky and J. Isaacson, *Phys. Rev. A* **20**, 2130 (1979).